

201 The Applicability of Using Straight Ureteral Stents for the Treatment of Ureteral Stones in Presumably Non-Compliant Patients

Mydlo, J.H.; Streater, S. (Brooklyn, N.Y.)

205 Intracellular Electrical Activity in Human Urinary Bladder Smooth Muscle: The Effect of High Sucrose Medium

Visser, A.J.; van Maastrigt, R. (Rotterdam)

Case Reports

212 Prostatic Malacoplakia Associated with Prostatic Abscess: Diagnosis and Treatment

Arena, F. (Parma); Fortunati, C. (Borgo val di Taro); di Stefano, C.; Peracchia, G.; Cortellini, P. (Parma)

216 Primary Squamous Cell Carcinoma of the Prostate: A Rare Clinicopathological Entity. Report of 2 Cases and Review of Literature

Nabi, G.; Ansari, M.S.; Iqbal Singh, I.; Sharma, M.C.; Dogra, P.N. (New Delhi)

220 MALT-Type Primary Lymphoma of the Urinary Bladder: Clinicopathological Study of 2 Cases and Review of the Literature

Wazait, H.D.; Chahal, R.; Sundurum, S.K.; Rajkumar, G.N.; Wright, D.; Aslam, M.M. (West Yorkshire)

225 An Unusual Cause of Irritable Urinary Bladder Symptoms

Reisman, Y.; de Reijke, T.M. (Amsterdam)

227 Transvesical Placement of One Limb of an Aortobifemoral Bypass Graft as a Complication of Aortic Bypass Surgery

Farkas, P.; Lüdecke, G.; Akintürk, H.; Weidner, W. (Giessen)

229 Handicap of Walking by a Huge Paratesticular Liposarcoma

Augustin, H.; Schips, L.; Vilitis, P.; Zigeuner, R.; Petritsch, H.P.; Uggowitzler, M.; Hubmer, G. (Graz)

196 Erratum

219 Ankündigung

231 Author Index Vol. 66, 2001

233 Subject Index Vol. 66, 2001

S. Karger
Medical and Scientific Publishers
Basel • Freiburg • Paris • London
New York • New Delhi • Bangkok
Singapore • Tokyo • Sydney

Drug Dosage

The authors and the publisher have exerted every effort to ensure that drug selection and dosage set forth in this text are in accord with current recommendations and practice at the time of publication. However, in view of ongoing research, changes in government regulations, and the constant flow of information relating to drug therapy and drug reactions, the reader is urged to check the package insert for each drug for any change in indications and dosage and for added warnings and precautions. This is particularly important when the recommended agent is a new and/or infrequently employed drug.

All rights reserved.

No part of this publication may be translated into other languages, reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, recording, microcopying, or by any information storage and retrieval system, without permission in writing from the publisher or, in the case of photocopying, direct payment of a specified fee to the Copyright Clearance Center (see 'General Information').

© Copyright 2001 by S. Karger AG,
P.O. Box, CH-4009 Basel (Switzerland)
Printed in Switzerland on acid-free paper by
Reinhardt Druck, Basel