

- 4 Swindle P, Eastham JA, Ohori M, Kattan MW, Wheeler T, Maru N, et al: Do margins matter? The prognostic significance of positive surgical margins in radical prostatectomy specimens. *J Urol* 2008;179(5 suppl):S47–S51.
- 5 Meeks JJ, Eastham JA: Radical prostatectomy: positive surgical margins matter. *Urol Oncol* 2013;31:974–979.
- 6 Oncoline: Richtlijn Prostaatacarcinoom 2.0. [www.oncoline.nl](http://www.oncoline.nl). Integraal Kankercentrum Nederland, 2014.
- 7 Eastham JA, Kattan MW, Riedel E, Begg CB, Wheeler TM, Gerigk C, et al: Variations among individual surgeons in the rate of positive surgical margins in radical prostatectomy specimens. *J Urol* 2003;170:2292–2295.
- 8 Karakiewicz PI, Eastham JA, Graefen M, Cagiannos I, Stricker PD, Klein E, et al: Prognostic impact of positive surgical margins in surgically treated prostate cancer: multi-institutional assessment of 5831 patients. *Urology* 2005;66:1245–1250.
- 9 Yamamoto S, Yonese J, Kawakami S, Ohkubo Y, Tatokoro M, Komai Y, et al: Preoperative serum testosterone level as an independent predictor of treatment failure following radical prostatectomy. *Eur Urol* 2007;52:696–701.
- 10 Namura K, Onuki T, Kouno M, Ishigaki H, Kato Y, Tajiri T, et al: [Retrospective analysis of predicting biochemical recurrence after initial radical prostatectomy]. *Hinyokika Kiyo* 2013;59:485–489.
- 11 Cary KC, Paciorek A, Fuldeore MJ, Carroll PR, Cooperberg MR: Temporal trends and predictors of salvage cancer treatment after failure following radical prostatectomy or radiation therapy: an analysis from the CaPSURE registry. *Cancer* 2014;120:507–512.
- 12 Eastham JA, Kuroiwa K, Ohori M, Serio AM, Gorbonos A, Maru N, et al: Prognostic significance of location of positive margins in radical prostatectomy specimens. *Urology* 2007;70:965–969.
- 13 Graefen M, Noldus J, Pichlmeier U, Haese A, Hammerer P, Fernandez S, et al: Early prostate-specific antigen relapse after radical retropubic prostatectomy: prediction on the basis of preoperative and postoperative tumor characteristics. *Eur Urol* 1999;36:21–30.
- 14 Simon MA, Kim S, Soloway MS: Prostate specific antigen recurrence rates are low after radical retropubic prostatectomy and positive margins. *J Urol* 2006;175:140–144; discussion 144–145.
- 15 Marks RA, Koch MO, Lopez-Beltran A, Montironi R, Juliar BE, Cheng L: The relationship between the extent of surgical margin positivity and prostate specific antigen recurrence in radical prostatectomy specimens. *Hum Pathol* 2007;38:1207–1211.
- 16 Sofer M, Hamilton-Nelson KL, Civantos F, Soloway MS: Positive surgical margins after radical retropubic prostatectomy: the influence of site and number on progression. *J Urol* 2002;167:2453–2456.
- 17 Pettus JA, Weight CJ, Thompson CJ, Middleton RG, Stephenson RA: Biochemical failure in men following radical retropubic prostatectomy: impact of surgical margin status and location. *J Urol* 2004;172:129–132.
- 18 Richters A, Derks J, Husson O, Van Onna IE, Fossion LM, Kil PJ, et al: Effect of surgical margin status after radical prostatectomy on health-related quality of life and illness perception in patients with prostate cancer. *Urol Oncol* 2015;33:16.e9–e15.
- 19 Preston MA, Breau RH, Lantz AG, Morash C, Gerridzen RG, Doucette S, et al: The association between nerve sparing and a positive surgical margin during radical prostatectomy. *Urol Oncol* 2015;33:18.e1–e6.
- 20 Bolla M, van Poppel H, Tombal B, Vekemans K, Da Pozzo L, de Reijke TM, et al: Postoperative radiotherapy after radical prostatectomy for high-risk prostate cancer: long-term results of a randomised controlled trial (EORTC trial 22911). *Lancet* 2012;380:2018–2027.

## Erratum

In the article by Tauber et al. [*Urol Int* 2015;95:324–328], entitled ‘Efficacy management of urolithiasis: flexible ureteroscopy versus extracorporeal shockwave lithotripsy’, the fourth author’s first name has been misspelled: Wolfgang Schimetta should read Wolfgang Schimetta.